

3D geological model of the Paleogene deposits in bauxitebearing district Snižnica (Posušje, BiH): from visualization to finding new bauxite deposits

Pavičić, Ivica; Dragičević, Ivan; Urumović, Kosta; Grubišić, Ivan

Source / Izvornik: **Abstracts book / 36th International Meeting of Sedimentology, 2023, 549 - 549**

Conference paper / Rad u zborniku

Publication status / Verzija rada: **Published version / Objavljena verzija rada (izdavačev PDF)**

Permanent link / Trajna poveznica: <https://urn.nsk.hr/urn:nbn:hr:169:801338>

Rights / Prava: [In copyright](#) / [Zaštićeno autorskim pravom.](#)

Download date / Datum preuzimanja: **2025-03-15**



Repository / Repozitorij:

[Faculty of Mining, Geology and Petroleum Engineering Repository, University of Zagreb](#)





36TH



IAS

DU
BROV
NIK

MEETING OF SEDIMENTOLOGY

ABSTRACTS BOOK



12-16 June 2023, DUBROVNIK, CROATIA

36th International Meeting of Sedimentology
June 12–16, 2023, Dubrovnik, Croatia

ABSTRACTS BOOK



Organized by:

Croatian Geological Society (HGD) and International Association of Sedimentologists (IAS)



Organizing Committee

Lara Wacha, **chair**, *Croatian Geological Survey, Zagreb*
Katarina Gobo, *University of Zagreb, Faculty of Science*
Nikolina Ilijanić, *Croatian Geological Survey, Zagreb*
Tvrтко Korbar, *Croatian Geological Survey, Zagreb*
Marijan Kovačić, *University of Zagreb, Faculty of Science*
Duje Kukoč, *Croatian Geological Survey, Zagreb*
Borna Lužar-Oberiter, *University of Zagreb, Faculty of Science*
Maja Martinuš, *University of Zagreb, Faculty of Science*
Slobodan Miko, *Croatian Geological Survey, Zagreb*
Davor Pavelić, *University of Zagreb, Faculty of Mining, Geology and Petroleum Engineering*
Kristina Pikelj, *University of Zagreb, Faculty of Science*
Igor Vlahović, *University of Zagreb, Faculty of Mining, Geology and Petroleum Engineering*

Scientific Committee

Igor Vlahović, **president**, *University of Zagreb, Croatia*
Nevena Andrić Tomašević, *Karlsruhe Institute of Technology, Germany*
Bruno Campo, *University of Bologna, Italy*
Sonia Campos Soto, *Complutense University of Madrid, Spain*
Luca Caracciolo, *FAU Erlangen-Nürnberg, Germany*
Blanka Cvetko Tešović, *University of Zagreb, Croatia*
Shahin E. Dashtgard, *Simon Fraser University, Canada*
Andrea Di Capua, *National Research Council – IGAG, Italy*
Goran Durn, *University of Zagreb, Croatia*
Gianluca Frijia, *University of Ferrara, Italy*
Massimiliano Ghinassi, *University of Padova, Italy*
Luis Gibert Beotas, *University of Barcelona, Spain*
Bosiljka Glumac, *Smith College, USA*
Antun Husinec, *St. Lawrence University, USA*
Stuart Jones, *Durham University, UK*
Tvrтко Korbar, *Croatian Geological Survey, Croatia*
Marijan Kovačić, *University of Zagreb, Croatia*
Juan Carlos Laya, *Texas A&M University, USA*
Marta Marchegiano, *University of Granada, Spain*
Cole McCormick, *Pennsylvania State University, USA*
Mardi McNeil, *Geoscience Australia, Australia*
Theresa Nohl, *University of Vienna, Austria*
Shuxin Pan, *PetroChina – NWGI, China*
Guido Pastore, *University of Milano–Bicocca, Italy*
Maximiliano Paz, *University of Saskatchewan, Canada*
Daniel A. Petráš, *Czech Geological Survey, Czech Republic*
Miquel Poyatos-Moré, *Universitat Autònoma of Barcelona, Spain*
Joanna Pszonka, *Polish Academy of Sciences – MEERI, Poland*
John J.G. Reijmer, *Vrije Universiteit Amsterdam, The Netherlands*
Valentina Marzia Rossi, *National Research Council – IGG, Italy*
Arnoud Sloopman, *Colorado School of Mines, USA*
Miroslaw Slowakiewicz, *University of Warsaw, Poland*
Thomas Steuber, *Khalifa University of Science and Technology, Abu Dhabi, UAE*
Finn Surlyk, *University of Copenhagen, Denmark*
Michal Šujan, *Comenius University in Bratislava, Slovakia*
Romain Vaucher, *University of Geneva, Switzerland*
Alan Vranjković, *INA Oil Company, Croatia*
Lara Wacha, *Croatian Geological Survey, Croatia*
Guodong Wang, *PetroChina, China*
Pujun Wang, *Jilin University, China*
Valentin Zuchuat, *RWTH Aachen University, Germany*
Nadja Zupan Hajna, *Research Centre of the Slovenian Academy of Sciences and Arts, Slovenia*

Publisher: Croatian Geological Society (HGD)

For the publisher: Slobodan Miko

Editors: Igor Vlahović and Darko Matešić

Language Editor: Julie Robson (Scotland, United Kingdom)

Digital layout: Laser Plus d.o.o

Cover design: Ana Badrić

eISBN: 978-953-6907-79-3

Theme 16. Techniques and technologies in sedimentology**General Session**

Poster presentation

3D Geological model of the Paleogene deposits in bauxite-bearing district Snižnica (Posušje, BiH): from visualization to finding new bauxite deposits

Ivica Pavičić¹, Ivan Dragičević¹, Kosta Urumović², Ivan Grubišić³¹University of Zagreb, Faculty of Mining, Geology and Petroleum Engineering, Zagreb, Croatia²Croatian Geological Survey, Department of Hydrogeology and Engineering Geology Zagreb, Croatia³Rudnici boksita d.o.o. Posušje, Posušje, Bosnia and Herzegovinaivica.pavicic@rgn.hr

One of the economically most valuable bauxite deposits in the Adriatic Carbonate Platform (AdCP) originates from the terrestrial phase between the Upper Cretaceous and Paleogene. The bauxites formed during Upper Cretaceous–Paleogene emersion are present on the entire AdCP. One of the largest and most explored and exploited area is the bauxite-bearing area of Posušje in Bosnia and Herzegovina. Geological exploration and exploitation were continuous and intensive between 1950 and 1990, but after that, all works were stopped by war. In the last ten years, research has started again, and exploitation has been stable with a tendency to increase. This work results from the new research combined with old data collected by Rudnici Boksita d.o.o. Posušje. Footwall to the bauxite deposits are thick Cretaceous rudist limestones, and the hanging-wall is composed of diverse carbonate and clastic sediments of the Paleogene age. This work aimed to collect all available data, systematize them into a 3D geodatabase and construct a 3D geological model of Paleogene strata in Snižnica locality, one of the most complex districts in the area. This work results in a new geological map, 15 geological cross-sections, and a 3D geological model of the Paleogene strata. The 3D model shows the spatial distribution of different Paleogene lithofacies, and the thickness of Paleogene deposits, which is important for drilling planning and reconstruction of eroded parts of the Paleogene deposits. The thickness of Paleogene deposits ranges from 0 to almost 200 m, while the eroded part of Paleogene deposits reaches up to 100 m. The 3D geological model also indicates that the largest amount and largest deposits are located under the clastic series of deposits, especially if the eroded part is taken into account. The constructed 3D geological model and the conducted analysis can serve as a basis for planning further exploration works, primarily drill-holes to find new bauxite deposits.