

Determining the formation of kaolinite mineralization using mineralogical analyses in Motajica granite complex, Bosnia and Herzegovina

Brenko, Tomislav; Borojević Šoštarić, Sibila

Source / Izvornik: **Knjiga sažetaka = Book of abstracts / 7. hrvatski geološki kongres s međunarodnim sudjelovanjem, 2023, 33 - 34**

Conference paper / Rad u zborniku

Publication status / Verzija rada: **Published version / Objavljena verzija rada (izdavačev PDF)**

Permanent link / Trajna poveznica: <https://urn.nsk.hr/urn:nbn:hr:169:895922>

Rights / Prava: [In copyright](#) / [Zaštićeno autorskim pravom.](#)

Download date / Datum preuzimanja: **2024-07-12**



Repository / Repozitorij:

[Faculty of Mining, Geology and Petroleum Engineering Repository, University of Zagreb](#)



prior to testing the algorithms, the data were standardized, normalized, or transformed depending on the algorithms used (SEMENIKHIN & BELOZEROV, 2019).

Based on the correlation matrix, 14 variables (logs) were determined for the calculation of the forecast model. A total of six regression models were used to predict the acoustic logs and porosity values. The base model was calculated using the Support Vector Regression (SVR) algorithm (Fig. 1), as the goal was to determine the boundaries of acceptable predictions.

The measure of accuracy for regression models, amongst others, is the coefficient of determination R^2 , which indicates how well the predicted and measured val-

ues correlate. For the base model, at the shallower depths the predicted values follow the trend of the measured ones but at the greater depths the predictions fall out of the trend, which makes the results unreliable. The most accurate model was generated using the K-Nearest Neighbour (KNN) algorithm, where the coefficient of determination was over 0.9 (Fig. 1).

The greatest advantages of predicting the values using trained models are the ability to fill in missing logging data and the short time required to get the initial image of the subsurface. Based on the predicted values, the further processing and interpretation methods are determined more quickly.

KRPAN, B., PLEIĆ, M., TAKAČ, D., BIGUNAC, D., MATKOVIĆ, M., KRIZMANIĆ, K., MAVAR, A., SULAIMAN, W., PERICA, R., MIŠUR, A., TROSKOT ČORBIĆ, T., DEVIĆ, V., ČORKOVIĆ, D. (2018): Regionalna studija za istražni prostor DR-02. Fond tehničke dokumentacije INA – Industrija nafte d.d., 203 p.

MCDONALD, A. (2021): Data Quality Considerations for Petrophysical Machine Learning Models. SPWLA 62nd Annual Logging Symposium, Society of Petrophysicists and Well Log Analysts, 1–25.

SEMENIKHIN, A., BELOZEROV, B.V. (2019): Well Log Data Standardization, Imputation and Anomaly Detection Using Hidden Markov Models. Fourth EAGE Conference on Petroleum Geostatistics, EAGE, 1–4.

TADEJ, J. (2011): Razvoj ranomiocenskih i srednjomiocenskih taložnih okoliša sjeverozapadnog dijela Dravske depresije na temelju podataka iz dubokih bušotina. Doktorski rad. Rudarsko-geološko-naftni fakultet, Zagreb, 168 p.

DETERMINING THE FORMATION OF KAOLINITE MINERALIZATION USING MINERALOGICAL ANALYSES IN MOTAJICA GRANITE COMPLEX, BOSNIA AND HERZEGOVINA

UTVRĐIVANJE MEHANIZMA NASTANKA KAOLINITA KORIŠTENJEM MINERALOŠKIH ANALIZA U MOTAJIČKOM GRANITNOM KOMPLEKSU, BOSNA I HERCEGOVINA

Tomislav Brenko¹, Sibila Borojević Šoštarić^{1*}

¹University of Zagreb, Faculty of Mining, Geology and Petroleum Engineering, Pierottijeva 6, 10 000 Zagreb, Croatia

*corresponding author: sibila.borojevic-sostaric@rgn.unizg.hr

Keywords: *kaolinitisation, hydrothermal alteration, XRD, FTIR*

The aim of this abstract is to present the formation mechanism of the kaolinite and kaolin deposits in the Motajica Mt. Although several projects and investments are currently being carried out in the Motajica Mt. area, the development and formation of kaolinite mineralisation is still relatively poorly known. Kaolinite deposits are associated with the Motajica granite pluton, which is characterised by syncollisional granite of Eocene age (JURKOVIĆ, 2004). Numerous papers have been published on the Motajica kaolinites (STANGAČILOVIĆ, 1956; JURKOVIĆ & ŠINKOVEC, 1960; VARIĆAK, 1966; TRUBELJA & BARIĆ, 1979), with different authors proposing different formation mechanisms for the kaolinites. Some authors believed that kaolinitisation occurred during Tertiary ma-

rine transgression, others suggested that kaolinitisation occurred through superficial weathering, while in some cases it was considered to be the result of hydrothermal solutions.

The focus of this work is mineralogical analysis of the kaolinitised granites in the Motajica area aiming at determination of clay minerals and definition of the kaolinite deposits origin. The following analyses were performed on the samples: macroscopic determination of mineral composition, X-ray diffraction analysis on original samples, samples treatment with ethylene glycol, heating at 350 °C and 550 °C and treatment with dimethyl sulphoxide (DMSO), Fourier transformed infrared spectroscopy (FTIR) and geochemical analysis on three samples.

The semi-quantitative mineral composition of the kaolinitised granites of the original samples revealed typi-

cal granite mineralisation, including quartz, K-feldspars, plagioclase and mica minerals. Diffractograms obtained after treating the samples with different treatment methods show a relatively strong diffraction peak of illite at 10 Å and a diffraction peak of kaolinite at 7 Å in all samples (Fig. 1). In addition, some samples indicate chlorite, possibly due to alteration of biotite granite. Faint indicative peaks of topaz and dickite were also noted. The presence of these minerals suggests that hydrothermal activity in the temperature range 100–350 °C played an important role in the formation of the kaolinite mineralisation. FTIR spectrum indicates characteristic absorption bands of kaolinite between 3500 and 3700 cm⁻¹. When four characteristic bands (approx. 3700, 3670, 3650 and 3620 cm⁻¹) are well defined, the structure of kaolinite is considered to be ordered (TIRONI *et al.*, 2012). Kaolinitised samples exhibit slightly higher Al contents than the locally occurring various granite types. Based on TiO₂/Zr and Cr+Nb/Ti+Fe ratios, two samples are consistent with purely hydrothermal kaolinitisation, while one sample shows signs of mixed type of formation (DILL *et al.*, 1997).

This hypothesis is consistent with the geochemical analysis of quartz veins occurring in the Bosanski Kobaš kaolin deposit in the north-eastern part of Motajica. Elevated contents of bismuth (Bi) in locally occurring quartz veins, similar to the results of RAMOVIĆ *et al.* (1982),




Figure 1. X-ray diffractogram of one Motajica sample with sample treatments

indicate strong hydrothermal activity in this area. The presence of hydrothermal veins in the samples studied clearly indicates a hydrothermal phase that played an important role in the formation of the kaolinites studied. Furthermore, several studies (FLEURENCE & NICOLAS, 1964; ESTÉOULE-CHOUX, 1981; SHOVAL *et al.* 1999) confirmed that hydrothermal kaolinites tend to have higher crystallinity when compared to supergene kaolinites. The presence of a supergene phase of kaolinitisation cannot be completely ruled out because the deposits are at the surface, but it can be safely assumed that there was initial hydrothermal kaolinitisation.

DILL, H., BOSSE, H.-R., HENNING, K.H., FRICKE, A. (1997): Mineralogical and chemical variations in hypogene and supergene kaolin deposits in a mobile fold belt the Central Andes of northwestern Peru. *Mineralium Deposita*, 32, 149–163.

ESTÉOULE-CHOUX, J. (1981): Etude en microscopie électronique à balayage de quelques kaolins d'origines différentes: Apports de cette technique pour la compréhension de leurs génèses. *Clay Minerals*, 16, 279–288.

FLEURENCE, A., NICOLAS, J. (1964): Observations sur la notion d'ordre et de désordre de certains minéraux du groupe de la kaolinite. *Bulletin du Groupe Français Des Argiles*, 14, 149–162.

JURKOVIĆ, I., ŠINKOVEC, B. (1960): Proračun rezervi kaolina u Motajici. *Fond Geol zavoda, Sarajevo*.

RAMOVIĆ, E. (1982): Metalogenetska rajonizacija područja sjeveroistočne i dijela sjeverne Bosne. *Geol. Glasnik*, 27, 117–153.

SHOVAL, S., YARIV, S., MICHAELIAN, K.H., BOUDEULLE, M., PANCZER, G. (1999): Hydroxyl-stretching bands 'A' and 'Z' in Raman and infrared spectra of kaolinites. *Clay Minerals*, 34, 551–563.

STANGAČILOVIĆ, D. (1956): Motajički primarni kaolin. *Glasnik Prirodnjačkog muzeja srpske zemlje, verzija A*, 7/1, 1–11.

TRUBELJA, F., BARIĆ, IJ. (1979): Minerali Bosne i Hercegovine, knjiga I (Silikati). *Zemaljski muzej BiH u Sarajevu*, 1–452.

VARIČAK, D. (1966): Petrološka studija motajičkog granitnog masiva. *Posebno izdanje Geol. glasnika*, 9, 1–170.