

Occurrence and mineral chemistry of the titanian clinohumite from marble of Zorovac creek, Moslavačka Gora, Croatia

Garašić, Vesnica

Source / Izvornik: **Knjiga sažetaka = Book of abstracts / 7. hrvatski geološki kongres s međunarodnim sudjelovanjem, 2023, 57 - 58**

Conference paper / Rad u zborniku

Publication status / Verzija rada: **Published version / Objavljena verzija rada (izdavačev PDF)**

Permanent link / Trajna poveznica: <https://urn.nsk.hr/urn:nbn:hr:169:776058>

Rights / Prava: [In copyright](#) / [Zaštićeno autorskim pravom.](#)

Download date / Datum preuzimanja: **2025-03-01**



Repository / Repozitorij:

[Faculty of Mining, Geology and Petroleum Engineering Repository, University of Zagreb](#)



OCCURRENCE AND MINERAL CHEMISTRY OF THE TITANIAN CLINOHUMITE FROM MARBLE OF ZOROVAC CREEK, MOSLAVAČKA GORA, CROATIA

POJAVA I MINERALNA KEMIJA TITANSKOG KLINOHUMITA IZ MRAMORA POTOKA ZOROVAC, MOSLAVAČKA GORA, HRVATSKA

Vesnica Garašić^{1*}

¹University of Zagreb, Faculty of Mining, Geology and Petroleum Engineering, Pierottijeva 6, 10 000 Zagreb, Croatia

*corresponding author: vesnica.garasic@rgn.unizg.hr

Keywords: *titanian clinohumite, marble, Moslavačka Gora*

Clinohumite is a scarce mineral, but is the most abundant mineral of the humite group which is characterized by a general formula of $nM_2SiO_4 \cdot xM_{1-x}Ti_x(OH, F)_2 \cdot 2xO_2 \cdot x$, where M stays for octahedrally coordinated Mg, Fe, Mn, Ca and Zn cations in declining order of abundance, $x < 1$ with $n=1, 2, 3, 4$ for norbergite, chondrodite, humite and clinohumite (JONES *et al.*, 1969). The occurrence of clinohumite is in the nature limited to only five rock types: kimberlites, Archean ultramafics, Alpine peridotites, carbonatites and marbles. However, despite the rarity of its occurrence, clinohumite is essential in understanding hydration and dehydration processes in the petrogenesis of mantle derived ultramafic bodies found in polymetamorphic Archean terranes (NISHIO *et al.*, 2019), but also in reconstruction of Ti mobility in the frame of crustal metamorphism linked to the clinohumite formation in forsterite marbles (KARMAKAR, 2021). Recently, the occurrence of humite-bearing marbles has been used for the reconstruction of the plate geometries during Neoproterozoic time, known as “humite epoch”, characterized by very water-rich, fluorinated fluid activity coeval with the waning stages of Pan-African tectono-thermal event (PRADEEPKUMAR & KRISHNANATH, 2000; FERNANDES & CHAVES, 2014).

Clinohumite occurs in Croatia only in the grey marbles of Zorovac creek in the Moslavačka Gora and was described by BARIĆ (1972), GARAŠIĆ (1993) and BALEN *et al.* (2000). The Moslavačka gora is crystalline complex located in the SW part of the Pannonian Basin, in Croatia. It comprises metamorphic rocks of high-to medium-grade, predominately migmatites and gneisses and of medium grade, mostly metapelites and amphibolites, which intermittently surround granitoids of different kind (PAMIĆ, 1990). The Moslavačka Gora belongs to Sava zone (SCHMID *et al.*, 2008), located between Laurasia and Gondwana, and was recognised as high-heat flow zone causing Cretaceous igneous and metamorphic events (BALEN & PETRINEC, 2011). Zircon dating of two mica-granite gave Cretaceous age and Early Ordovician age for metagranite (the most of the metamorphic complex), whereas the Cretaceous age of low pressure-high temperature (LP/HT) metapelite was obtained by monazite dating (STARIJAS *et al.*, 2010).

The studied marbles occur as interlayers up to 50 cm thick in cordierite and diopside-amphibole schists. Mineral assemblages in marbles indicate three metamorphic events of different metamorphic grade. The highest grade mineral assemblage consists of forsterite+spinel+calcite+pargasite. The most important forsterite+clinohumite+calcite±phlogopite±tremolite assemblage occurred during LP/HT metamorphism, as result of replacement of forsterite by clinohumite. The textural relationships in the lowest grade mineral assemblage consisting of clinohumite+spinel+calcite+chlorite+dolomite suggest that chlorite and dolomite grew at the expense of clinohumite, spinel and calcite (Fig. 1). Also replacement of forsterite by talc and serpentine belongs to this retrograde lowest grade metamorphism.




Figure 1. The lowest grade mineral assemblage in marbles of Moslavačka Gora consisting of clinohumite (Chu), spinel (Sp), calcite (Cc), chlorite (Chl) and dolomite (Do)

Microprobe analyses of clinohumite revealed that content of TiO₂ varies between 0.88 (in the mineral parageneses with phlogopite) and 4.04 wt.% (in the mineral parageneses with amphiboles). The fluorine concentration in Ti-clinohumite shows negative correlation with titanium, and ranges from 1.30 to 3.16 wt.%. This is in accordance with expected substitution (Mg,Fe)+2(OH,F) = Ti+2(O). The ratio of F/(F+OH) in clinohumite or X_F^{Chu} is in the range between 0.32 (in the parageneses with amphiboles) and 0.57 (in the parageneses with phlogopite). Content of H₂O varies from 2.16 to 2.66 wt.%. The concentrations of MgO reach the values from 51.87 to 55.63 wt.% depending on the content of titanium and iron oxides, whereas FeO content ranges from 1.64 to 4.79 wt.%. The SiO₂ content shows slight variation (37.07 and 37.63 wt.%). Other cations in Ti-clinohumite are present in negligible amounts.

Textural relationships and microprobe analyses reveal that Ti-clinohumite grew at the expense of forsterite caused by infiltration of titanian and fluorinated H₂O-rich fluids. Such rare clinohumite marbles with remarkably similar mineral assemblages, textures and P-T-fluid metamorphic conditions and multiphase evolution are widespread in the dispersed Gondwana fragments (PRADEEPKUMAR & KRISHNANATH, 2000). Although the Cretaceous age of LP/HT metapelite was determined in the Moslavačka Gora, because marbles are more sensitive than other rocks to the changes in fluid composition (FERRY, 1992), the LP/HT metamorphism in marbles is not necessarily comparable to the LP/HT metamorphism in the metapelites.

I thank Rainer Altherr for his help in using the electron microprobe at the Institut für Petrographie und Geochemie, Fridericiana Universität Karlsruhe (TH), Germany.

- BALEN, D., BELAK, M., TIBLJAŠ, D., TOMAŠIĆ, N. (2000): The succession of metamorphic parageneses in mineral assemblage from marble – Zorovac Creek (Moslavačka gora, Northern Croatia). Second Croatian Geological Congress, Proceedings, 93–96.
- BALEN, D., PETRINEC, Z. (2011): Contrasting tourmaline types from peraluminous granites: a case study from Moslavačka Gora (Croatia). *Mineralogy and petrology*, 102, 117–134.
- BARIĆ, LJ. (1972): Kontaktnometamorfni mramori iz okolice Podgarića u Moslavačkoj gori (Hrvatska). VII kongres geologija SFRJ, Predavanja II knjiga, 121–137.
- FERNANDES, M.L.S., CHAVES, A.O. (2014): Chemical composition and genesis of the clinohumites from marbles of Itaoca-Girona, Espirito Santo State, Brazil. *Comunicacoes Geologicas*, 101, 81–84.
- FERRY, J.M. (1991): Dehydration and decarbonation reactions as a record of fluid infiltration. In: Kerrick, D.D. (ed.): *Contact metamorphism*, Reviews in Mineralogy, 26, Mineral. Soc. Amer., 26, 351–393.
- GARAŠIĆ, V. (1993): Uvjeti metamorfizma stijena amfibolitnog facijesa Moslavačke gore. Unpublished M.Sc. Thesis, University of Zagreb, 150 p.
- JONES, N.W., RIBBE, P.H., GIBBS, G.V. (1969): Crystal chemistry of humite minerals. *American Mineralogist*, 54, 391–411.
- KARMAKAR, S. (2021): Formation of clinohumite±spinel in dolomitic marbles from the Makrohar Granulite Belt, Central India: Evidence for Ti mobility during regional metamorphism. *American Mineralogist*, 106, 1818–1827.
- NISHIO, I., MORISHITA, T., SZILAS, K., PEARSON, G., TANI, K.I., TAMURA, A., HARIGANE, Y., GUOTANA, J. (2019): Titanian clinohumite-bearing peridotite from the Ulamertoq ultramafic body in the 3.0 Ga Akia Terrane of Southern West Greenland. *Geosciences*, 9, 153.
- PAMIĆ, J. (1990): Alpine granites, migmatites and metamorphic rocks from Mt. Moslavačka Gora and the surrounding basement of the Pannonian Basin (Northern Croatia, Yugoslavia). *Rad Jugoslavenske akademije znanosti i umjetnosti Zagreb*, 10, 7–121.
- PRADEEPKUMAR, A.P., KRISHNANATH, R. (2000): A Pan-African “humite epoch” in East Gondwana: Implications for Neoproterozoic Gondwana geometry. *Journal of Geodynamics*, 29, 43–62.
- SCHMID S.M., BERNOULLI, D., FÜGENSCHUH, B., MATENCO, L., SCHEFFERR, S., SCHUSTER, R., TISCHLER, M., USTASZEWSKI, K. (2008): The Alpine-Carpathian-Dinaridic orogenic system: correlation and evolution of tectonic units. *Swiss J. Geosci.*, 11, 139–183.
- STARIJAJŠ, B., GERDES, A., BALEN, D., TIBLJAŠ, D., FINGER, F. (2010): The Moslavačka Gora crystalline massif in Croatia: A Cretaceous heat dome within remnant Ordovician granitoid crust. *Swiss J. Geosci.*, 103, 61–82.