

Geomorphological settings and types of landslides in the city of Buzet identified using LiDAR digital terrain model

Jagodnik, Petra; Bernat Gazibara, Sanja; Sinčić, Marko; Lukačić, Hrvoje; Arbanas, Željko; Mihalić Arbanas, Snježana

Source / Izvornik: **Knjiga sažetaka = Book of abstracts / 7. hrvatski geološki kongres s međunarodnim sudjelovanjem, 2023, 79 - 79**

Conference paper / Rad u zborniku

Publication status / Verzija rada: **Published version / Objavljena verzija rada (izdavačev PDF)**

Permanent link / Trajna poveznica: <https://um.nsk.hr/um:nbn:hr:169:170999>

Rights / Prava: [In copyright](#) / [Zaštićeno autorskim pravom.](#)

Download date / Datum preuzimanja: **2025-03-16**



Repository / Repozitorij:

[Faculty of Mining, Geology and Petroleum Engineering Repository, University of Zagreb](#)



GEOMORPHOLOGICAL SETTINGS AND TYPES OF LANDSLIDES IN THE CITY OF BUZET IDENTIFIED USING LIDAR DIGITAL TERRAIN MODEL

GEOMORFOLOŠKI OKOLIŠI I TIPOVI KLIZIŠTA NA PODRUČJU GRADA BUZETA IDENTIFICIRANI PRIMJENOM LIDAR DIGITALNOGA MODELA RELJEFA

Petra Jagodnik^{1*}, Sanja Bernat Gazibara², Marko Sinčić², Hrvoje Lukačić², Željko Arbanas¹, Snježana Mihalić Arbanas²

¹University of Rijeka, Faculty of Civil Engineering, Radmile Matejčić 3, 51 000 Rijeka, Croatia

²University of Zagreb, Faculty of Mining, Geology and Petroleum Engineering, Pierottijeva 6, 10 000 Zagreb

*corresponding author: petra.jagodnik@gradri.uniri.hr

Keywords: *visual interpretation of LiDAR DTM, landslide inventory map, geomorphological settings of landslides, landslides in gullies, Buzet*

This study presents the results of landslide detection and mapping at a large scale, performed in the area of the city of Buzet in central Istria. The research was conducted within the frame of scientific research project “Methodology development for landslide susceptibility assessment for land-use planning based on LiDAR technology” (LandSlidePlan, HRZZ IP-2019-04-9900). The study area (20 km²) consists of Eocene flysch sediments, composed of a rhythmical alternation of marl and carbonate sediments in the lower part of the complex, and thinly bedded carbonate-siliciclastic turbidite sediments in the upper part (BERGANT *et al.*, 2003).

For the first time, the detailed geomorphological landslide inventory map (GUZZETTI *et al.*, 2012) is created for the area in the flysch environment in central Istria, using innovative remote sensing technology that proved to be effective in mapping landslides (GÖRÜM, 2019; JAGODNIK *et al.*, 2020a). Identification and mapping of landslides was carried out based on the visual interpretation of topographic datasets derived from the bare-earth LiDAR (Light Detection and Ranging) Digital Terrain Model (DTM) at a 0.3 m spatial resolution. Airborne laser scanning was performed in March 2020, with an average point density of 16 points per m².

In the study area, more than 1,160 landslides are identified and delineated with high geographical accuracy and thematic certainty due to the clear visibility of landslide features on LiDAR DTM derivatives. However, it was quite a challenge to identify and map individual landslides in areas of gully erosion and badlands, which represent the typical geomorphological phenomena in the flysch environment of central Istria (GULAM *et al.*, 2014). Landslide density is 58 landslides per km². Most of the landslides are debris slides, and debris slide-debris flows (HUNGR *et al.*, 2014), which are the main types of landslides in flysch deposits (JAGODNIK *et al.*, 2020b). Landslides are predominantly small and shallow. Their sizes are in the range between only 4 m² to 8 ha.

Generally, there are three typical geomorphological settings of landslides in the study area: (i) complex gullies; (ii) agricultural fields; and (iii) artificial slopes along the roads. Gullies are the predominant environment for the occurrence of landslides, with approximately 65 % of identified landslides being situated in gullies. Such specific geomorphological setting of landslide phenomena confirmed that there is a significant interplay between mass movements and fluvial processes in the investigated area. Therefore, in the future research, the results of this study will be used for testing the relevance of gully and badland phenomena as conditioning factors in landslide susceptibility modelling at a large scale.

BERGANT, S., TIŠLIJAR, J., ŠPARICA, M. (2003): Eocene carbonates and flysch deposits of the Pazin basin. In: Vlahović, I., Tišljarić, J. (eds.), 22nd IAS meeting of sedimentology – Opatija 2003. Excursion Guide-Book, Institute of Geology, Zagreb, 57–63.

GÖRÜM, T. (2019): Landslide recognition and mapping in a mixed forest environment from airborne LiDAR data. *Engineering Geology*, 258, 105155.

GULAM, V., POLLAK, D., PODOLSKA, L. (2014): The analysis of the flysch badlands inventory in central Istria, Croatia. *Geologia Croatica*, 67/1, 1–15.

GUZZETTI, F., MONDINI, A.C., CARDINALI, M., FIORUCCI, F., SANTANGELO, M., CHANG, K.T. (2012):

Landslide inventory maps: new tools for an old problem. *Earth-Science Reviews*, 112, 42–66.

HUNGR, O., LEROUEIL, S., PICARELLI, L. (2014): The Varnes classification of landslide types, an update. *Landslides*, 11, 167–194.

JAGODNIK, P., JAGODNIK, V., ARBANAS, Ž., MIHALIĆ ARBANAS, S. (2020a): Landslide types in the Slani Potok gully. *Geologia Croatica*, 73/1, 13–28.

JAGODNIK, P., BERNAT GAZIBARA, S., ARBANAS, Ž., MIHALIĆ ARBANAS, S. (2020b): Engineering geological mapping using airborne LiDAR datasets – an example from the Vinodol Valley, Croatia. *Journal of Maps*, 16/2, 856–867.