

Genesis and epigenetic evolution of the Minjera bauxites, Croatia, Istria

Perković, Ivor; Durn, Goran; Matešić, Darko; Vlahović, Igor; Cvetko Tešović, Blanka; Martinuš, Maja

Source / Izvornik: Knjiga sažetaka = Book of abstracts / 7. hrvatski geološki kongres s međunarodnim sudjelovanjem, 2023, 141 - 142

Conference paper / Rad u zborniku

Publication status / Verzija rada: Published version / Objavljena verzija rada (izdavačev PDF)

Permanent link / Trajna poveznica: <https://urn.nsk.hr/urn:nbn:hr:169:135221>

Rights / Prava: [In copyright](#) / Zaštićeno autorskim pravom.

Download date / Datum preuzimanja: **2024-05-08**



Repository / Repozitorij:

[Faculty of Mining, Geology and Petroleum Engineering Repository, University of Zagreb](#)



ljena biometrijska analiza kućica ove vrste (CORDEY *et al.*, 1970), stoga su procedura i odabir mjerjenih parametara uglavnom slijedili metodologiju opisanu tim radom. 50 kućica vrste *P. micra* nasumično je odabrano i fotografirano s bočne i spiralne strane. Svakoj kućici izmjereno je najveći promjer, visina i širina najmlađe klijetke, te broj klijetaka u zadnjem zavoju. Uz to identificiran je tip ušća, koje kod ove vrste varira od visokog luka do bipartitnog, podijeljenog na dva luka.

Mjerenjem je utvrđeno da u svim istraženim uzorcima prosjek maksimalnog promjera kućice iznosi od 140,3 do 159,9 µm, što je manje od holotipa (170 µm). Najmanji maksimalni promjer kućice imaju u uzorku iz zone E15, u starijem dijelu naslaga uvale Zaraće, a to je ujedno i

uzorak u kojem ova vrsta ima najveću zastupljenost u zajednici (11,3 %). Širina i visina najmlade klijetke variraju neovisno o ostalim mjerjenim parametrima dok je prosječan broj klijetaka u zadnjem zavoju ujednačen, krećući se između 6,2 i 6,4. Bipartitan tip ušća je vrlo rijedak, te niti u jednom uzorku broj jedinki s ovakvim ušćem ne prelazi 4 %. Ovi rezultati ukazuju da je *P. micra* u sutropskim do umjerenim uvjetima kakvi su vladali tijekom gornjeg eocena na prostoru današnjeg otoka Hvara, imala kućice manjih dimenzija od pripadnika iste vrste iz tropskih područja.

Istraživanje je financirala Hrvatska zaklada za Znanost u sklopu projekta BREEMECO (IP-2019-04-5775).

- CORDEY, W.G., BERGGREN, W.A., OLSSON, R.K. (1970): Phylogenetic Trends in the Planktonic Foraminiferal Genus *Pseudohastigerina* Banner and Blow, 1959. *Micropaleontology*, 16/2, 235–242.
- LUCIANI, V., GIUSBERTI, L., AGNINI, C., FORNACIARI, E., RIO, D., SPOFFORTH, D.J.A., PÄLKE, H. (2010): Ecological and evolutionary response of Tethyan planktonic foraminifera to the middle Eocene climatic optimum (MECO) from the Alano section (NE Italy). *Palaeogeography, Palaeoclimatology, Palaeoecology*, 292, 82–95.

- PEARSON, P.N., OLSSON, R.K., SPEZZAFERI, S., LECKER, I.E. (2018): Taxonomy, biostratigraphy and phylogeny of Oligocene Globanomalinidae (*Pseudohastigerina* and *Turborotalia*). In: Wade, B.S., Olsson, R.K., Pearson, P.N., Huber, B.T., Berggren, W.A. (eds.), *Atlas of Oligocene planktonic foraminifera*, Cushman foundation for foraminiferal research, 524 p.
- WADE, B.S., PEARSON, P.N., BERGGREN, W.A., PÄLKE, H. (2011): Review and revision of Cenozoic tropical planktonic foraminiferal biostratigraphy and calibration to the geomagnetic polarity and astronomical time scale. *Earth-Science Reviews*, 104, 111–142.

GENESIS AND EPIGENETIC EVOLUTION OF THE MINJERA BAUXITES, CROATIA, ISTRIA GENEZA I EPIGENETSKI RAZVOJ BOKSITA MINJERE, HRVATSKA, ISTRA

Ivor Perković^{1*}, Goran Durn¹, Darko Matešić¹, Igor Vlahović¹, Blanka Cvetko Tešović², Maja Martinuš²

¹University of Zagreb, Faculty of Mining, Geology and Petroleum Engineering, Pierottijeva 6, 10 000 Zagreb, Croatia

²University of Zagreb, Faculty of Science, Department of Geology, Horvatovac 102b, 10 000 Zagreb, Croatia

*corresponding author: ivor.perkovic@rgn.unizg.hr

Keywords: grey and pyritised bauxite, morphology of iron sulphides, influence of marine porewater, palaeotopographical positions

Pyritisation of bauxites occurs as a result of their epigenetic reduction during the formation of swampy environments in the bauxite cover during the initial phase of the transgression, which commonly follows the formation of the bauxite. The Minjera bauxites, which formed during the terrestrial phase which lasted between the Late Cenomanian/Late Santonian and Early Eocene, were pyritised. The epigenetic pyritisation of these bauxites was related to the transgression that followed their formation, which led to the formation of ponds and swamps in the paleo-depressions in the karstified terrain, such as the sinkholes and can-

yons filled with the bauxite. Therefore, all of the Minjera bauxite bodies are covered by Lower Eocene Liburnian beds formed in such environments.

The Minjera bauxites have been mined in the past, but only the pyrite-containing bauxite was used in the production of alum and vitriol, while the red bauxite was left in the tailing heaps in the area. Pyritised bauxite samples from two Minjera bauxite bodies (D-1 and D-15) were collected, with the aim to reconstruct their genesis and the processes which led to their formation and subsequent pyritisation. Two main types of bauxites, the grey bauxite and the pyritised bauxite, were distinguished based on their mineralogy, geochemistry as well as their structure and texture. The grey bauxite contains

high amounts of kaolinite and moderate to high amounts of diaspore, while containing no or little böehmite. Iron sulphides, represented mainly by pyrite and sporadically by marcasite, are generally present in very low amounts in this type of bauxite and appear as veinlets and crystal clusters in the matrix and bauxite clasts and as replacements of iron oxide rich laminae in ooids. This type of bauxite is also enriched in bases and large ion lithophile elements compared to pyritised bauxite, which is likely related to lower leaching intensity. Pyritised bauxite contains high amounts of böehmite and iron sulphides, while containing very little to moderate amounts of kaolinite, and almost no diaspore. Iron sulphides appear in these samples in many different morphologies and textures. They replace the iron oxide-rich laminae within the ooids and the fine-grained matrix between the bauxite clasts and ooids. In samples where the matrix was not completely pyritised, the pyritisation started from many crystallisation centres, from which iron sulphides grew outward, either in the forms of frambooids or rosettes composed of needle shaped crystals. Iron sulphides in these samples commonly crystallize along the fractures, which is also seen on a large scale, in the form of centimetre-thick veins of iron sulphides parallel with the boundary between the bedrock and the bauxite. Both bauxites are enriched in heavy REEs, and display a slight negative cerium anomaly, which indicates the influence of marine porewater. Different textures and morphologies of iron sulphides suggest variations in the saturation with

iron and sulphur and are probably linked with the sea-level variations during the initial stages of the transgression, which could have affected the production of organic matter in swampy environments that developed in the cover of the bauxite during this stage. The ingress of marine porewater was most likely the source of sulphur, which was derived from the microbial reduction of sulphur in the organic matter-rich environment. The pyritisation appears to have affected each bauxite deposit differently, since the grey bauxite is almost exclusively found in the D-15 deposit, while the pyritised bauxite is found only in the D-1 deposit. Besides their distinct epigenetic evolution, this also suggests that bauxites of different grades formed contemporaneously in the area, as the D-15 body is composed mainly from highly kaolinitic grey bauxite and the D-1 body from the highly böehmitic pyritised bauxite. This is probably related to different morphologies of the two bauxite bodies, as the D-1 body is much larger and steeper (> 20 m) than the D-15 bauxite body (< 5 m). The differences in their morphologies likely developed as a consequence of their different palaeotopographical positions, which led to different rates of chemical weathering between the two bauxite bodies.

This work has been fully supported by Croatian Science Foundation under the project IP-2019-04-8054 – WianLab (Western Istrian Anticline as an Ideal Natural Laboratory for the Study of the Regional Unconformities in Carbonate Rocks).

NON-KARST AREAS IN THE KARST AREA OF CROATIA NEKRŠKA PODRUČJA NA PODRUČJU KRŠA RH

Lea Petohleb^{1*}, Nikolina Bakšić-Pavlović¹, Ivona Žiža¹, Stjepan Husnjak², Vladimir Kušan¹

¹Oikon d.o.o. Institut za primijenjenu ekologiju, Trg Senjskih uskoka 1-2, 10 020 Zagreb Hrvatska

²Agronomski fakultet, Sveučilište u Zagrebu, Svetosimunska cesta 25, 10 000 Zagreb, Hrvatska

*dopisni autor: lpetohleb@oikon.hr

Ključne riječi: krš, GIS, poljoprivredna područja, ruralni razvoj, rekultivacija

Za potrebe Programa ruralnog razvoja Republike Hrvatske (RH) za programsko razdoblje 2021. – 2027. godine izrađen je projekt Određivanje područja s posebnim ograničenjima ili ostalim posebnim ograničenjima s kalkulacijama uz utvrđivanje vrijednosti kontekst indikatora broj 41 „Organska tvar u tlu“ i broj 42 „Erozija tla vodom“ za programsko razdoblje 2021. – 2027. Uredbom (EU) br. 1305/2013, propisano je da ukupna površina područja s posebnim (specifičnim) ograničenjima može iznositi maksimalno 10 % u odnosu na ukupnu površinu države, a što bi u slučaju Hrvatske iznosilo 5659,4 km².

U skladu s time RH je predložila da se u područje s posebnim (specifičnim) ograničenjima uvrste otoci i poluotok Pelješac. Osim toga, predloženo je da se u područje s posebnim (specifičnim) ograničenjima uvrsti i prostor krša u kojem su osim depopulacije i zapuštanja poljoprivrede prisutni i vrlo teški uvjeti za bavljenje poljoprivrednom proizvodnjom uvjetovani vrlo razvedenim krškim reljefom. Prije svega, u području krša, poljoprivredna proizvodnja je otežana zbog velikog broja vrtaca po jedinici površine i vrlo često velikom kamenitošću površine.

Krško područje prikazano je i analizirano prema definiciji krša iz Izvješća o stanju u prostoru RH (2003).