

Late Aptian and Early Albian regional emersion phase in Istria, Croatia

Mileusnić, Marta; Durn, Goran; Ottner, Franz; Fröschl, Heinz; Tišljar, Josip

Conference presentation / Izlaganje na skupu

Permanent link / Trajna poveznica: <https://um.nsk.hr/um:nbn:hr:169:425391>

Rights / Prava: [In copyright](#)/[Zaštićeno autorskim pravom.](#)

Download date / Datum preuzimanja: **2025-03-17**



Repository / Repozitorij:

[Faculty of Mining, Geology and Petroleum Engineering Repository, University of Zagreb](#)






7th International Symposium on the Cretaceous
5-9 September 2005

Scientific Program and Abstracts

Conveners: Karl B. Föllmi and Thierry Adatte



Edited by Alexis Godet, Haydon Mort, Pascal Linder and Stéphane Bodin



during this important stage of their evolution.

similar to Purbeckian sediments of NW Europe.

LATE APTIAN AND EARLY ALBIAN REGIONAL EMERSION PHASE IN ISTRIA, CROATIA

Mileusnic Marta¹, Durn Goran¹, Ottner Franz²,
Fröschl Heinz³ and Tišljarić Josip¹

¹Faculty of mining, geology and petroleum engineering, University of Zagreb, Zagreb, HR 10000, Croatia; ²Department für Bautechnik und Naturgefahren, University of Natural Resources and Applied Life Sciences, Vienna, A 1190, Austria; ³Austrian Research Centers GmbH – ARC, Seibersdorf, A-2444, Austria

Emails: : marta.mileusnic@rgn.hr; gdurn@rgn.hr;
franz.ottnner@boku.ac.at; jtisljar@rgn.hr

Istrian part of the Adriatic carbonate platform emerged in the Middle/Late Aptian for the second time and was exposed for 11-19 MY. Exposure related features, which mark type 1 sequence boundary, are represented by greenish-grey clays, marls and lime breccias. Mineralogical, micromorphological and geochemical analysis were performed on clay layers, ranging in thickness from several centimetres up to 1 meter, and the insoluble residues of limestones situated immediately below clay layers.

The main mineral components of clays are illitic material and illite/smectite mixed layer minerals. Occasionally, pyrite, gypsum and chlorite are present. In contrast, insoluble residue is dominated by smectite and illitic material, and occasionally kaolinite. The clay mineral distribution of one investigated profile show a clear trend indicating the influence of both pedogenic and diagenetic processes. Micromorphological data support consideration of pedogenetic alteration. Distribution of vanadium, uranium, molybdenum and REE, as colour of clay and presence of abundance of pyrite framboids indicate unoxic environment. The current investigation should give answers to provenance of the material from which this clays are formed.

We consider that the clays are remnants of seasonally marshy soils to permanently waterlogged soil, which formed from the erosional remains of surficial soils and sediments, which were accumulated in palaeokarst pits following an oscillating marine transgression that terminated emergence. The Lower Cretaceous shallow-marine deposition sporadically interrupted by periods of emersion are sedimentologically and palaeogeographically very

OCEAN ANOXIC EVENTS IN THE MID-CRETACEOUS SIMULATED BY 3-D BIOGEOCHEMICAL GENERAL CIRCULATION MODEL

Misumi Kazuhiro¹, Yamanaka Yasuhiro¹

¹Graduate School of Environmental Earth Science,
Hokkaido University, Sapporo, 060-0810, Japan

Emails: : misumi@ees.hokudai.ac.jp, galapen@ees.hokudai.ac.jp

The mid-Cretaceous is well known by its Ocean Anoxic Event (OAEs). Decreased supply or increased consumption of dissolved O₂ in the deep water is regarded as a cause for the OAEs. But there is little consensus as to the overall forces behind the OAEs. In this study, we conducted a mid-Cretaceous simulation by an OGCM combined with the biogeochemical cycle model to investigate causes of OAEs. The model results show that the thermohaline circulation (THC) oscillated between activity and inactivity. A stable enlargement of an anoxic region is only seen in the proto-Atlantic. This is due to the high export production supported by the elevated concentration of PO₄, which prevents the supplying of dissolved O₂ to the deep water even under the active THC. Mechanisms that enhance the PO₄ concentration are needed to explain the large temporal and spatial scales that are associated with the OAEs.

AMMONITES FROM THE BASE OF THE RYAZANIAN STAGE AND BOREAL-TETHYAN CORRELATION

Mitta Vasily

Paleontological Institute of RAS, Moscow, 117997
Russia

Email: mita@paleo.ru

The Zone Riasanites rjasanensis is the lowest Zone of the Ryazanian Stage and Cretaceous System. This Zone is presented in Russian platform by strongly condensed phosphate sandstones by thickness up to 1-2 m with numerous fossils. The Rjasanensis Zone is characterized by two groups of ammonites which are quite different in the origin. They are "Boreal" Craspeditidae and "Tethyan" Neocomitidae. The

