

Late Albian silicification event(s) in Istria, Croatia

Barudžija, Uroš; Aljinović, Dunja; Mileusnić, Marta; Durn, Goran

Source / Izvornik: **Scientific Program and Abstracts, 2005**

Conference presentation / Izlaganje na skupu

Permanent link / Trajna poveznica: <https://urn.nsk.hr/urn:nbn:hr:169:974941>

Rights / Prava: [In copyright](#) / [Zaštićeno autorskim pravom.](#)

Download date / Datum preuzimanja: **2024-04-19**



Repository / Repozitorij:

[Faculty of Mining, Geology and Petroleum
Engineering Repository, University of Zagreb](#)





7th International Symposium on the Cretaceous

5-9 September 2005

Scientific Program and Abstracts

Conveners: Karl B. Föllmi and Thierry Adatte



Edited by Alexis Godet, Haydon Mort, Pascal Linder and Stéphane Bodin



LATE ALBIAN SILICIFICATION EVENT(S) IN ISTRIA, CROATIA

Barudzija Uros¹, Aljinovic Dunja¹, Mileusnic Marta¹
and Durn Goran¹

¹Faculty of Mining, Geology and Petroleum
Engineering, University of Zagreb, Zagreb, HR
10000, Croatia

Emails: ubarud@rgn.hr; daljin@rgn.hr;
marta.mileusnic@rgn.hr; gdurn@rgn.hr

In the NW part of the Adriatic Carbonate Platform (AdCP), in Istria region, during Late Albian diagenetic silicification occurred. Silicification took place within the peritidal carbonate sequences comprising intertidal carbonate breccias and laminated cyanobacterial limestones, as well as subtidal mudstones peloidal wackstones/packstones. Three silicified layers can be traced, more or less continuously, along the 30km traverse. According to their position in the sedimentary column and to their geometry, they are interpreted as three possible events of aeolian input of silica (volcanic ash) to carbonate platform environment. Sedimentological, mineralogical and geochemical analysis were performed on the samples from the silicified layers (ranging from 15cm to 2m in thickness), as well as from the related limestones and marls/clayey limestones. Silicification changed primary carbonate sediments in places almost completely and formed so-called "quartz diagenetic sediments" (up to 98% of SiO₂), which contain several types of quartz (idiomorphic authigenic quartz, microcrystalline quartz aggregates, chert-like clusters). Distributions of microelements and REE indicate differences of the silicified layers from surrounding limestones. Insoluble residues of clayey limestones predominantly contain illite-smectite minerals, together with quartz, some feldspars and amorphous material.

These results support assumption about the input of the source material for silicification to carbonate platform. Current investigations (SEM) should give additional informations about the source and mechanisms of silicification.

RECORD OF THE FARAONI ANOXIC EVENT (LATE HAUTERIVIAN) IN THE ULTRAHELVETIC DOMAIN

Baudin François¹, Busnardo Robert², Beltran
Catherine¹, de Rafelis Marc¹, Renard Maurice¹,
Charollais Jean³ and Clavel Bernard²

¹Département de Géologie Sédimentaire et CNRS-
FR 32, Université Pierre et Marie Curie, case 117,
4 place Jussieu 75252 Paris cedex 05, France;

²Département des Sciences de la Terre, Université de
Lyon, 27-43 bd du 11 Novembre 69622 Villeurbanne
Cedex, France; ³Département de Géologie et
Paléontologie, Université de Genève, 13 rue des
Maraîchers 1211 Genève, Switzerland

An equivalent of the Faraoni Level, originally described in the Umbria-Marche Basin (central Italy) as an upper Hauterivian regional marker-bed, and then identified in southern Alps (Italy), Vocontian Basin (SE France) and Subbetic Basin (Spain), was recently recognised in the Ultrahelvetetic domain along the Veveyse de Châtel-Saint-Denis section (external Prealps, Switzerland) by Busnardo *et al.* (2003).

The trend in the evolution of oxygen and carbon isotopic ratios and trace elements (Sr, Mn, Fe) from the carbonates show that a 4-meter thick critical interval of the upper Hauterivian (*Pseudothurmania angulicostata* ammonite Zone) succession of the Veveyse de Châtel-Saint-Denis section experienced strong environmental changes. This interval coincides with an increase in marine organic matter (1.5 % TOC and hydrogen index values up to 460 mg HC/g TOC).

This organic matter enrichment associated with changes in the ocean chemistry is probably the record of the Faraoni 'Anoxic Event', although its sedimentary expression in the Ultrahelvetetic domain is less obvious compared to other Tethyan basins (Baudin *et al.*, submitted). It should be noted, however, that the sequential position of this level is exactly the same in the different basins, e.g. the basal part of a transgressive phase.

This new site allows to extend the distribution of this late Hauterivian anoxic event in an intermediate paleogeographic domain between the Umbria-Marche and the Vocontian Basins.

References

Baudin F., Busnardo R., Beltran C., de Rafelis M., Renard M., Charollais J. & Clavel B. (submitted) - Enregistrement de l'événement anoxique Faraoni (Hauterivien supérieur) dans le domaine

