

# The landslide susceptibility assessment for application in the spatial planning system: from national to local scale

---

**Bernat Gazibara, Sanja; Sinčić, Marko; Krkač, Martin; Lukačić, Hrvoje; Šarić, Gabrijela; Jagodnik, Petra; Mihalić Arbanas, Snježana**

*Source / Izvornik:* **Knjiga sažetaka = Book of abstracts / 7. hrvatski geološki kongres s međunarodnim sudjelovanjem, 2023, 22 - 23**

**Conference paper / Rad u zborniku**

*Publication status / Verzija rada:* **Published version / Objavljena verzija rada (izdavačev PDF)**

*Permanent link / Trajna poveznica:* <https://um.nsk.hr/um:nbn:hr:169:259669>

*Rights / Prava:* [In copyright](#)/[Zaštićeno autorskim pravom.](#)

*Download date / Datum preuzimanja:* **2024-07-30**



*Repository / Repozitorij:*

[Faculty of Mining, Geology and Petroleum Engineering Repository, University of Zagreb](#)



# THE LANDSLIDE SUSCEPTIBILITY ASSESSMENT FOR APPLICATION IN THE SPATIAL PLANNING SYSTEM: FROM NATIONAL TO LOCAL SCALE

## PROCJENA PODLOŽNOSTI NA KLIZANJE ZA PRIMJENU U SUSTAVU PROSTORNOG PLANIRANJA: OD DRŽAVNOG DO LOKALNOG MJERILA

Sanja Bernat Gazibara<sup>1\*</sup>, Marko Sinčić<sup>1</sup>, Martin Krkač<sup>1</sup>, Hrvoje Lukačić<sup>1</sup>, Gabriijela Šarić<sup>1</sup>, Petra Jagodnik<sup>2</sup>, Snježana Mihalić Arbanas<sup>1</sup>

<sup>1</sup>University of Zagreb Faculty of Mining, Geology and Petroleum Engineering, Pierottijeva 6, 10 000 Zagreb, Croatia

<sup>2</sup>University of Rijeka, Faculty of Civil Engineering, Radmile Matejčić 3, 51 000 Rijeka, Croatia

\*corresponding author: sanja.bernat@rgn.unizg.hr

Keywords: *landslides, landslide susceptibility assessment, Croatia*

The main motivation to research the landslide susceptibility assessment for application in land use planning arises from the national landslide risk assessment (BERNAT GAZIBARA *et al.*, 2019) which recognised landslides as a second natural risk in Croatia (CNPDRR, 2019). Furthermore, the preliminary regional landslide susceptibility analysis showed that approx. 20 % of the Republic of Croatia area is potentially prone to sliding. Therefore, landslide susceptibility assessment for national, county and local levels was carried out in the frame of two scientific projects Methodology development for landslide susceptibility assessment for land-use planning based on LiDAR technology (LandSlidePlan, HRZZ IP-2019-04-9900) (BERNAT GAZIBARA *et al.*, 2022) and project Applied landslide research for the development of risk mitigation and prevention measures (PRI-MJER, KK.05.1.1.02.0020). The national landslide susceptibility map at a small scale is created to give a general overview of critical areas for an entire country, and its purpose is to inform policymakers and the general public (MIHALIĆ ARBANAS *et al.*, 2022). County-level landslide susceptibility assessment on a medium scale synthesizes available data and identifies wider areas with landslide problems and can be used to define areas for more detailed research on a local level. The third level is the local-scale landslide mapping and zonation that includes specific areas of local administrative units (municipality or city) or complex critical areas. The results were landslide susceptibility maps for seven study areas: (i) the Republic of Croatia; (ii) City of Zagreb, Karlovac County and Primorje–Gorski Kotar County; and (iii) the study areas in the Zagreb City (BERNAT GAZIBARA *et al.*, 2023), Hrvatsko Zagorje (SINČIĆ *et al.*, 2022a), Karlovac City (SINČIĆ *et al.*, 2022b) and Istria.

Methodology development for landslide susceptibility assessment on national and county scales was carried out using a heuristic approach, i.e. Fuzzy Logic method, and

available topographical and geological data. Given that the validation of the final landslide susceptibility map is mandatory, and systematic landslide inventories at the national or county level do not exist, we used the landslide database conducted by the University of Zagreb, Faculty of Mining, Geology and Petroleum Engineering. The database consists of 2,186 landslides with the exact location of the occurrence. All landslide susceptibility maps showed high accuracy and were classified into three susceptibility zones, considering The Area Under the Receiver Operating Characteristic Curve ( $AUC_{ROC}$ ).

Methodology development for landslide susceptibility assessment on a local scale was carried out using different mapping units and statistical methods (e.g. Information Value method, Weights of Evidence method, Logistic Regression and Discriminant Analysis, and machine learning methods, including Support Vector Machine, Artificial Neural Network and Random Forest). Moreover, landslide susceptibility models were computed using different scenarios of high-resolution input data, i.e. geometrical types of LiDAR-based inventory and variations of causal factors. Finally, all landslide susceptibility models were evaluated based on model fitting performance, model prediction performance, and model uncertainty. The purpose of comparing landslide susceptibility models is to define the most suitable maps for application in spatial planning at national, regional, and local levels. The research was based on innovative technologies, limitations related to the availability of spatial data in Croatia (limited amount of geological data), and urgent needs for efficient solutions applicable in the Croatian spatial planning system in line with European and global requirements related to sustainable development, human and environmental protection.

This work has been partially supported by University of Rijeka (project Uniri-tehnic 18-97 and Uniri-tehnic 18-298).

BERNAT GAZIBARA, S., CINDRIĆ KALIN, K., ERAK, M., KRKAČ, M., ĐOMLJJA, P. ARBANAS, Ž., MIHALIĆ ARBANAS, S. (2019): Landslide hazard analysis in national-scale for landslide risk assessment in Croatia. In: Ul-

jarević, M., Zekan, S., Salković, S., Ibrahimović, DŽ. (eds.): Proceedings of the 4th Regional Symposium on Landslides in the Adriatic-Balkan Region, Geotechnical Society of Bosnia and Herzegovina, 175–182.

- BERNAT GAZIBARA, S., MIHALIĆ ARBANAS, S., SINČIĆ, M., KRKAČ, M., LUKAČIĆ, H., JAGODNIK, P., ARBANAS, Ž. (2022): LandSlidePlan – Scientific research project on landslide susceptibility assessment in large scale. In: Peranić, J., Vivoda Prodan, M., Bernat Gazibara, S., Krkač, M., Mihalčić Arbanas, S., Arbanas, Ž. (eds.): *Landslide Modelling & Applications: Proceedings of the 5th Regional Symposium on Landslides in the Adriatic-Balkan Region*, Faculty of Civil Engineering, University of Rijeka and Faculty of Mining, Geology and Petroleum Engineering, University of Zagreb, 99–106.
- BERNAT GAZIBARA, S., SINČIĆ, M., KRKAČ, M., LUKAČIĆ, H., MIHALIĆ ARBANAS, S. (2023): Landslide susceptibility assessment on a large scale in the Podsljeme area, City of Zagreb (Croatia). *Journal of Maps*, 19, 1–11.
- CNPDRR, CROATIAN NATIONAL PLATFORM FOR DISASTER RISK REDUCTION (2019): *Disaster risk evaluation for The Republic of Croatia*, Zagreb, 180 p. (in Croatian)
- MIHALIĆ ARBANAS, S., BERNAT GAZIBARA, S., KRKAČ, M., SINČIĆ, M., LUKAČIĆ, H., JAGODNIK, P., ARBANAS, Ž. (2022): Landslide Detection and Spatial Prediction: Application of Data and Information from Landslide Maps. In: Alcantara-Ayala, I., Arbanas, Ž., Huntley, D., Konagai, K., Mikoš, M., Sassa, K., Sassa, S., Tang, H., Tiwari, B. (eds.): *Progress in Landslide Research and Technology*, Volume 1 Issue 2, Springer, 195–212.
- SINČIĆ, M., BERNAT GAZIBARA, S., KRKAČ, M., LUKAČIĆ, H., MIHALIĆ ARBANAS, S. (2022a): The Use of High-Resolution Remote Sensing Data in Preparation of Input Data for Large-Scale Landslide Hazard Assessments. *Land*, 11/8, 1–37.
- SINČIĆ, M., BERNAT GAZIBARA, S., KRKAČ, M., MIHALIĆ ARBANAS, S. (2022b): Landslide susceptibility assessment of the City of Karlovac using the bivariate statistical analysis. *Rudarsko-geološko-naftni zbornik*, 149–170.

## PETROGRAPHIC, MINERALOGICAL AND CHEMICAL CHARACTERISTICS OF BAUXITE FROM POSUŠJE DEPOSITS, BOSNIA AND HERZEGOVINA – AGEMERA PROJECT RESEARCH

### PETROGRAFSKE, MINERALOŠKE I KEMIJSKE KARAKTERISTIKE BOKSITA IZ LEŽIŠTA U POSUŠJU, BOSNA I HERCEGOVINA – AGEMERA PROJEKT ISTRAŽIVANJA

Šime Bilić<sup>1\*</sup>, Irena Peytcheva<sup>2</sup>, Stoyan Georgiev<sup>2</sup>, Atanas Hikov<sup>2</sup>, Ivica Pavičić<sup>1</sup>, Franjo Šumanovac<sup>1</sup>, Marko Holma<sup>3, 4, 5</sup>

<sup>1</sup> University of Zagreb, Faculty of Mining, Geology and Petroleum Engineering, Pierottijeva 6, 10 000 Zagreb, Croatia

<sup>2</sup> Bulgarian Academy of Sciences, Geological Institute, Acad. G. Bonchev, Bl. 24, 1113 Sofia, Bulgaria

<sup>3</sup> University of Oulu, Kerttu Saalasti Institute, PL 8000, 90 014 Oulu, Finland

<sup>4</sup> Muon Solutions Oy, Rakkarinne 9, 96 900 Saarenkylä, Finland

<sup>5</sup> Arctic Planetary Science Institute (APSI), Rakkarinne 9, 96 900 Saarenkylä, Finland

\*corresponding author: sime.bilic@rgn.hr

**Keywords:** *bauxite deposit, karst bauxite, Posušje, petrography, AGEMERA*

The exploitation of bauxite in the wider area of Posušje in Bosnia and Herzegovina has been known since the first half of the 20<sup>th</sup> century, and to date more than 1100 deposits or occurrences of various sizes and structures have been discovered. Bauxite deposits of Posušje represent typical karst bauxites, formed during the emersion phase between Upper Cretaceous rudist limestones in the footwall and various carbonate and clastic rocks of Paleocene/Eocene and Oligocene ages in the hanging wall. The most intensive mineralogical and geochemical bauxite research in these areas was carried out during the 70s and 80s. Afterwards, despite numerous newly discovered deposits, no petrographic, mineralogical and geochemical analysis on the bauxites has been published. The present study is focused on recently discovered bauxite deposits

in Posušje and their petrographic and mineralogical description. The bauxites show ooidic to pelitomorphic, but in places also clastic textures. Micro-ooids and spheroids are mostly smaller than 0.1 mm, and in places fragments of the former ooid can be found. In some bauxites, pebbles of resedimented bauxite prevail, mostly larger than 2 mm but sometimes larger than 1 cm, which define their conglomerate texture. Gibbsite and boehmite are the main Al-bearing minerals of the bauxites, and diasporite has not been observed. The ratio of gibbsite to boehmite varies between deposits, and in some deposits boehmite is the only aluminium phase present. Where present, the gibbsite is developed into relatively large hypidiomorphic crystals larger than 0.25 mm, with clearly defined polysynthetic lamellae. Hematite is the dominant Fe-phase in all deposits. Although rarely, goethite-rich bauxites can be found in the form of isolated layers, with just traces or no hematite. Zircon, apatite and calcite are minor and ac-